

DIABETES ACTION PLAN TYPE

School Settings

Use in conjunction with
Diabetes Management Plan.

This plan is not suitable for
Type 1 Diabetes

Insulin Injections

Photo	Child name
	Date of birth
	Grade / year

Name of school

Parent / carer name

Contact no.

Diabetes treating team

Hospital ur no.

Contact no.

Authorised by

Signature

Role

Date plan created

Plan does not expire.
Review is recommended in 12 months.

LOW Hypoglycaemia (hypo)

Blood Glucose Level (BGL) less than 4.0 mmol/L

Signs and symptoms:

Pale, headache, shaky, sweaty, dizzy, drowsy, changes in behaviour

Note:

Check BGL if hypo suspected. Symptoms may not always be obvious

- Do not leave child alone
- Do not delay treatment
- Treatment to occur where child is at time of hypo

• Hypo supplies located:

MILD*

Child conscious

(Able to eat hypo food) *mild is common

Step 1:

Give fast acting carbohydrate

Step 2:

Recheck BGL in 15 mins

- If BGL less than 4.0, repeat Step 1
- If BGL greater than or equal to 4.0, go to Step 3

Step 3:

Give slow acting carbohydrate

Step 3a:

If insulin is due & BGL greater than or equal to 4.0, give usual insulin dose & then eat meal immediately.

Step 4:

Resume usual activity when BGL 4.0 or higher

SEVERE

Child drowsy / unconscious

(Risk of choking / unable to swallow)

First Aid DRABCD

Stay with child

**CALL AN
AMBULANCE
DIAL 000**

Contact parent / carer when safe to do so

HIGH Hyperglycaemia (Hyper)

Blood Glucose Level (BGL) greater than or equal to 15.0 mmol/L is well above target and requires additional action

Signs and symptoms:

Increased thirst, extra toilet visits, poor concentration, irritability, tiredness

Note:

Symptoms may not always be obvious

Child well

- Encourage 1-2 glasses water per hour
- Return to usual activity
- Extra toilet visits may be required
- Re-check BGL in 2 hours

In 2 hours, if BGL still greater than or equal to 15.0,

**CALL PARENT/
CARER FOR
ADVICE**

Child unwell (e.g. vomiting)

- Contact parent/carers to collect child ASAP
- Check ketones

KETONES

If unable to contact parent/carers and blood ketones greater than or equal to 1.0 mmol/L.

**CALL AN
AMBULANCE
DIAL 000**

Use in conjunction with Diabetes Action Plan.

Tick boxes that apply

Insulin Administration

Insulin is given multiple times per day.

The child requires an injection of insulin:

- ☐ Before breakfast at before school care
- ☐ Lunchtime
- ☐ Other

Insulin injection minutes before a meal.

Carbohydrate food must always be eaten after a mealtime insulin injection.

The insulin dose for meals / snacks will be determined by:

- ☐ Set dose
- ☐ Flexible dosing guide/app
- ☐ Supervision required to ensure correct information added to app.

Location in the school where the injection is to be given:

Is supervision required? ☐ Yes ☐ No ☐ Remind only

Responsible staff will need training if they are required to:

- ☐ Administer injection (Dose as per additional documentation provided)
- ☐ Assist ☐ Observe
- ☐ Other diabetes medication required at school

See Medication Authority Form or relevant document

Disposal of medical waste

- Dispose of any used pen needles in sharps container provided.
- Dispose of blood glucose and ketone strips as per the school's medical waste policy.

Glucose Monitoring

Target range for glucose levels pre-meals: 4.0 – 7.0 mmol/L
7.1 – 14.9 mmol/L are outside target range requiring no action.

- Glucose levels outside the target range are common.
- A glucose check should occur where the child is at the time it is required

Blood Glucose Level (BGL) Monitoring – Fingerprick

(Used when a child is not wearing CGM or if the sensor falls out)

- Monitoring is performed using a fingerprick device and meter.
- Before doing a **blood glucose** check, the child should **wash and dry hands**.

Is the student able to do their own blood glucose level (BGL) check?

- ☐ Yes ☐ No (Support is required)

The responsible staff member needs to:

- ☐ Do the check ☐ Assist ☐ Observe ☐ Remind

Blood glucose levels (BGL) to be checked (tick all those that apply)

- ☐ Anytime hypo suspected ☐ Before snack ☐ Before lunch
☐ Before activity ☐ Before exams/tests ☐ When feeling unwell
☐ Beginning of after-school care session
☐ Other times – please specify

Continuous glucose monitoring (CGM)

- Continuous glucose monitoring consists of a small sensor that sits under the skin and measures glucose levels in the fluid surrounding the cells.
- If the sensor/transmitter falls out, staff to do BGL (Fingerprick) checks.
- A CGM reading can differ from a blood glucose level (BGL) reading during times of rapidly changing glucose levels e.g., eating, after insulin administration, during exercise.

A child wearing CGM must have a BGL check:

- ☐ Anytime hypo suspected. Hypo treatment is based on a BGL check
☐ When CGM reading less than mmol/L, must be confirmed. Follow Action Plan
☐ When CGM reading above mmol/L must be confirmed. Follow Action Plan
☐ When feeling unwell
☐ Sensor reading does not align with expectation or child's symptoms
☐ Other times – please specify

Low Blood Glucose Levels (Hypoglycaemia / Hypo)

FOLLOW ACTION PLAN

- If the child requires more than 2 consecutive fast acting carbohydrate treatments, as per their Diabetes Action Plan, call their parent/carer. Continue hypo treatment if needed while awaiting further advice.
- All hypo treatment should be provided by parent/carer.
- If the school is located more than 30 minutes from a reliable ambulance service, then staff should discuss Glucagon injection training with the child's Diabetes Treating Team.

Ketones

FOLLOW THE 'HYPERGLYCAEMIA ACTION PLAN'

- Ketones can be dangerous and occur most commonly in response to high glucose levels or if a child is unwell.

Eating and drinking

- If using flexible dosing all carbohydrate foods should be clearly labelled by the parent/carer with carbohydrate amounts in grams.
- It is not the responsibility of the school staff to count carbohydrates. However, school staff may need to assist a child to add up the carbohydrate amounts they wish to eat.
- Some children will need supervision to ensure all food is eaten.
- No food sharing.
- Seek parent/carer advice regarding foods for school parties/celebrations.
- Always allow access to water.

Name	
Hospital UR no.	
Date plan created	

Physical activity

Hypo treatment and a glucose monitoring device should always be with the child.

- Physical activity **may cause glucose levels to go high or low.**
- Some children may require a glucose level check before, during or after physical activity.
- Some children MAY require a slow acting carbohydrate before planned physical activity.

■ Additional information

Activity food

■ Activity food located

Glucose level range	Carbohydrate food	Amount

- Physical activity should not be undertaken if BGL less than 4.0 mmol/L.

REFER TO THE DIABETES ACTION PLAN FOR HYPO TREATMENT

- Physical activity **should not** be undertaken if the child is **unwell**.

Excursions / incursions

It is important to plan for extracurricular activities.

- Ensure blood glucose monitor, blood glucose strips, ketone strips, insulin pen and pen needles, hypo and activity food are readily accessible.
- Plan for meal and snack breaks.
- Always have hypo treatment available.
- Know location of toilets.

School camps

- Is there a school camp planned for this year? ☐ Yes ☐ No
- Parents/carers need to be informed of any school camp at least 2 months prior to ensure the child's diabetes treating team can provide a Camp Diabetes Management plan and any training needs required.
- A Camp Diabetes Management Plan is different to the usual School Plan.
- Parents/carers will need a copy of the camp menu and activity schedule.
- At least 2 responsible staff attending the camp require training to be able to support the child on camp.
- If the camp location is more than 30 minutes from a reliable ambulance service Glucagon injection training will be required.

Exams

- Glucose level should be checked and documented before an exam.
- Glucose level should be greater than 4.0 mmol/L before exam is started.
- Blood glucose monitor and blood glucose strips, CGM devices or smart phones, hypo treatments, and water should be available in the exam setting.
- Extra time will be required if a hypo occurs, for toilet privileges, or child unwell.

Applications for special consideration

National Assessment Program Literacy and Numeracy (NAPLAN)

Applies to Grade 3, Grade 5, Year 7, Year 9. Check National Assessment Program website – Adjustment for student with disability for further information.

Victorian Certificate of Education (VCE)

Should be lodged at the beginning of Year 11 and 12. Check Victorian Curriculum and Assessment Authority (VCAA) requirements.

Name	
Hospital UR no.	
Date plan created	

Equipment checklist

Supplied by the parent/carer.

- ☐ Insulin pens and pen needles. Stored according to the school Medication Policy.
- ☐ Finger prick device
- ☐ Blood glucose monitor
- ☐ Blood glucose strips
- ☐ Blood ketone strips
- ☐ Hypo treatment
- ☐ Activity food
- ☐ Sharps' container
- ☐ Charging cables for diabetes management devices
- ☐ Smart phone to be used as medical device

Equipment checklist

Glossary of terms

AGREEMENTS

Parent/Carer

Organise a meeting with the school representatives to discuss implementation and sign off on your child's action and management plan.

- ☐ I have read, understood, and agree with this plan.
- ☐ I give consent to the school to communicate with the Diabetes Treating Team about my child's diabetes management at school.

Name

First name (please print)

Family name (please print)

Signature

Date

School representative

- ☐ I have read, understood, and agree with this plan.

Name

First name (please print)

Family name (please print)

Role

☐ Principal

☐ Vice principal

Signature

Date

Diabetes Treating Medical Team

Name

First name (please print)

Family name (please print)

Signature

Date

Hospital name

Name	
Hospital UR no.	
Date plan created	